



GB04/3527



INVESTOR IN PEOPLE

# **PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

REC'D 20 SEP 2004

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

*Stephen Hordley*

Dated 3 September 2004



20.4.2004 14:00 COPY

Patents Form 1/77

Patents Act 1977  
(Rule 16)



19NOV03 E853142-1 D00192  
P01/7700 0.00-0326848.9

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your reference	P.89234A GCW		
2. Patent application number (The Patent Office will fill this part in)	18 NOV 2003	0326848.9	
3. Full name, address and postcode of the or of each applicant (underline all surnames)	MARS, INCORPORATED 6885 Elm Street McLean, Virginia 22101-3883 United States of America		
Patents ADP number (if you know it)			
If the applicant is a corporate body, give the country/state of its incorporation	Virginia, USA	7495039001	
4. Title of the invention	ALLELIC VARIANTS		
5. Name of your agent (if you have one)	J. A. KEMP & CO.		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	14 South Square Gray's Inn London WC1R 5JJ		
Patents ADP number (if you know it)	26001		
6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months.	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. Divisionals, etc: Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)	Number of earlier UK application		Date of filing (day / month / year)
8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request?	Yes		
Answer YES if:			
a) any applicant named in part 3 is not an inventor, or			
b) there is an inventor who is not named as an applicant, or			
c) any named applicant is a corporate body.			
Otherwise answer NO (See note d)			

Patents Form 1/77

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheets of this form

Description	47	↘
Claim(s)	5	↘
Abstract	1	↘
Drawing(s)	4 + 4	↘

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for a preliminary examination and search (Patents Form 9/77)

Request for a substantive examination (Patents Form 10/77)

Any other documents (please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature(s)

*J. A. Kemp & Co.*  
J.A. KEMP & CO.

Date 18 November 2003

12. Name, daytime telephone number and e-mail address, if any, of person to contact in the United Kingdom

WOODS, Geoffrey Corlett  
020 7405 3292

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered YES in part 8, a Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.

## ALLELIC VARIANTS

### Field of the invention

The present invention relates to the diagnosis and treatment of IgA-related disorders in animals, such as gastrointestinal, skin and respiratory disease, and to novel polynucleotides and polypeptides.

### Background of the invention

Dogs of the German shepherd breed are particularly susceptible to a number of inflammatory and immune-mediated alimentary diseases, including small intestinal bacterial overgrowth and inflammatory bowel disease (IBD). German shepherd dogs (GSD) have also been reported to have a relative IgA deficiency, on the basis of reduced concentrations of serum IgA as compared to control populations. Similar differences in IgA concentration have been reported in tears, duodenal juice, duodenal explant culture media and faeces of this breed when compared with other breeds of dog. A reduced concentration of IgA has been found in the duodenal juice from GSD with small intestinal bacterial overgrowth compared with normal dogs, and tissue culture supernatants from twenty-four hour duodenal explants from GSD with chronic diarrhoea produce less IgA than explants derived from affected dogs of other breeds. However, in the same populations there were no differences in the numbers of IgA<sup>+</sup> plasma cells or CD4<sup>+</sup> T-cells within the duodenal lamina propria.

IgA plays an important role in the immune defence of mucosal sites, where it is secreted at concentrations far in excess of other immunoglobulin classes. In this context, IgA prevents colonisation and invasion by microorganisms, neutralises bacterial toxins and is involved in the elimination of antigen from the subepithelial lamina propria. The IgA molecule is of particular relevance in the immunological defence of the gastrointestinal tract, and dysfunction of mucosal immunity contributes to a range of idiopathic inflammatory bowel diseases that occur in man and other species. Multiple IgA subclasses have been identified in humans, primates and lagomorphs, whereas mice, cattle and dogs have only a single subclass. The two human subclasses (IgA<sub>1</sub> and IgA<sub>2</sub>) are defined by a difference in the length of the hinge region between the CH<sub>1</sub> and CH<sub>2</sub> domains. The single IgA subclass identified in dogs has a hinge region with a predicted amino acid sequence similar to the IgA<sub>1</sub> subclass of humans.

### Summary of the invention

The present inventors have shown that dogs possess multiple IgA allelic variants. These variants differ principally within the 5' end of the second exon of the  $\alpha$  heavy chain gene, an area corresponding to the hinge region of the molecule. The presence of an extended hinge region of IgA makes it more susceptible to cleavage by proteases. Therefore differences in hinge length and composition between variants have significant effects on the function of the IgA molecules they encode. These differences result in certain genotypes having an increased susceptibility to disease caused by pathogens that produce proteases that can cleave the hinge region.

Accordingly, the invention provides a method for determining susceptibility to an IgA-related disorder in an animal, the method comprising:

- a) identifying the or each IgA allelic variant present in an animal; and
- b) thereby determining whether the animal is susceptible to an IgA-related disorder. The invention further provides:

- a probe, primer or antibody which is capable of detecting an IgA allelic variant;

- a kit for carrying out the method of the invention comprising means for detecting an IgA allelic variant;

- a method of preparing customised food for an animal which is susceptible to an IgA-related disorder, the method comprising:

- (a) determining whether the animal is susceptible to an IgA-related disorder by a method of the invention; and

- (b) preparing food suitable for the animal;

- a method of providing a customised animal food, comprising providing food suitable for an animal which is susceptible to an IgA-related disorder to the animal, the animal's owner or the person responsible for feeding the animal, wherein the animal has been genetically determined to be susceptible to an IgA-related disorder;

- a method for identifying an agent for the treatment of an IgA-related disorder, the method comprising:

- (a) contacting an IgA allelic variant polypeptide or a polynucleotide which encodes an IgA allelic variant with a test agent; and

- (b) determining whether the agent is capable of binding to the polypeptide or modulating the activity or expression of the polypeptide or polynucleotide;

- use of a compound which is therapeutic for an IgA-related disorder in the

manufacture of a medicament for the prevention or treatment of an IgA-related disorder in an animal that has been identified as being susceptible to an IgA-related disorder by a method of the invention;

- a method of treating an animal for an IgA-related disorder, the method comprising administering to the animal an effective amount of a therapeutic compound which prevents or treats the disorder, wherein the animal has been identified as being susceptible to an IgA-related disorder by a method of the invention;

- a database comprising information relating to IgA allelic variants and optionally their association with IgA-related disorder(s);

10 - a method for determining whether an animal is susceptible to an IgA-related disorder, the method comprising:

(a) inputting data of one or more IgA allelic variant(s) present in the animal to a computer system;

(b) comparing the data to a computer database, which database comprises information relating to IgA allelic variants and the IgA-related disorder susceptibility associated with the variants; and

(c) determining on the basis of the comparison whether the animal is susceptible to an IgA-related disorder;

- a computer program comprising program code means that, when executed on a computer system, instructs the computer system to perform a method according to the invention;

- a computer program product comprising a computer-readable storage medium having recorded thereon a computer program according to the invention;

- a computer program product comprising program code means on a carrier wave, which program code means, when executed on a computer system, instruct the computer system to perform a method according to the invention;

- a computer system arranged to perform a method according to the invention comprising:

(a) means for receiving data of the one or more IgA allelic variant(s) present in the animal;

(b) a module for comparing the data with a database comprising information relating to IgA allelic variants and the IgA-related disorder susceptibility associated with the variants; and

(c) means for determining on the basis of said comparison whether the animal is susceptible to an IgA-related disorder;

- a method of preparing customised food for an animal which is susceptible to an IgA-related disorder, the method comprising:

5 (a) determining whether the animal is susceptible to an IgA-related disorder by a method according to claim 23;

(b) electronically generating a customised animal food formulation suitable for the animal;

10 (c) generating electronic manufacturing instructions to control the operation of food manufacturing apparatus in accordance with the customised animal food formulation; and

(d) manufacturing the customised animal food according to the electronic manufacturing instructions;

15 - use of a computer system of the invention to make a customised animal food product.

- an isolated polynucleotide comprising:

(a) an IgA variant sequence that differs to SEQ ID NO: 1 at one or more polymorphic positions as defined herein;

(b) any one of SEQ ID NO:s 3, 5, 7 or 9;

20 (c) a sequence that is complementary or is degenerate as a result of the genetic code to a sequence as defined in (a) or (b); or

(d) a fragment of (a), (b) or (c) which differs to SEQ ID NO: 1 at one or more polymorphic positions as defined in claim 5 and which is at least 10 nucleotides in length; and

25 - a polypeptide comprising:

(a) a sequence encoded by a polynucleotide of the invention;

(b) any one of SEQ ID NO:s 4, 6, 8 or 10; or

30 (c) a fragment of (a) or (b) which differs to SEQ ID NO: 2 at one or more polymorphic positions as defined in claim 5 and which is at least 10 amino acids in length.

### **Brief description of the sequences**

SEQ ID NO: 1 shows the polynucleotide sequence of the canine IgA  $\alpha$ -chain, starting from the beginning of exon 1. SEQ ID NO: 2 shows the corresponding polypeptide sequence.

5 SEQ ID NO:s 3 and 4 show the polynucleotide and polypeptide sequences of variant A of the canine IgA  $\alpha$ -chain.

SEQ ID NO:s 5 and 6 show the polynucleotide and polypeptide sequences of variant B of the canine IgA  $\alpha$ -chain.

10 SEQ ID NO:s 7 and 8 show the polynucleotide and polypeptide sequences of variant C of the canine IgA  $\alpha$ -chain.

SEQ ID NO:s 9 and 10 show the polynucleotide and polypeptide sequences of variant D of the canine IgA  $\alpha$ -chain.

SEQ ID NO:s 11 to 17 show primer and probe sequences.

15 SEQ ID NO:18 shows the Genbank sequence L36871 (canine IgA  $\alpha$ -chain polynucleotide sequence).

### **Brief description of the drawings**

Figure 1 shows real-time RT-PCR quantification of mRNA expression in duodenal biopsies. The graph on the left (A) demonstrates the traces produced using primer set 1 showing the bimodal distribution of the expression. Samples either  
20 contained a relatively 'high' (●) or 'low' (■) amounts of  $\alpha$ -chain mRNA. The graph on the right shows the overlap of the traces when primer set 2 is used indicating that similar amounts of  $\alpha$ -chain mRNA is present in all samples but the forward primer in set 1 does not detect a significant portion of the mRNA in some samples.

25 Figure 2 shows the four sequenced variants and their relationship to the Genbank sequence L36871. The numbering starts from the first base of exon 1 of the Genbank sequence. A single nucleotide polymorphism (C or T) (A) exists between individuals with the same variant which does not alter the predicted amino acid sequence. A variable number of CT repeats (D) are present in the intron sequence prior  
30 to the splice site of the second exon. A single base polymorphism at base 547 (T or A) (E) causes loss of the splice acceptor resulting in either long hinge variants (F to H) or short hinge variants (G to H). The first exon ends at base 306 (B) and it corresponds to



the first domain of the heavy chain. Four bases in the intron sequence of all dogs sequenced do not agree with that of the Genbank sequence (C).

Figure 3 shows the mRNA sequences for the IgA variants together with the positions of the primer sets used in the real-time PCR. The positions are numbered from the first base of exon one of the Genbank sequence. The position of the forward primer in set one includes the 9 base addition of variant C and D and would therefore not efficiently prime off this template. The forward primer in set 2 can amplify fragments from all variants. The difference in length of the RNA can be seen, with variant B the shortest and D the longest.

Figure 4 shows the predicted amino acid sequence for the IgA variants based on the sequenced mRNAs. Position A corresponds to the (A-T) polymorphism in the first exon which has no effect on the encoded amino acid. The amino acid differences are highlighted by bold text. Variant B has the shortest hinge (7 AA) and variant D the longest (13AA).

Figure 5 shows an alignment of mRNA encoding human IgA<sub>1</sub> and IgA<sub>2</sub>. The mRNA sequences encoding the human IgA subclasses are aligned to demonstrate the position of the 39bp deletion of IgA<sub>2</sub> which results in the shorter hinge of this subclass. In contrast to the hinge variants of dogs, this deletion does include the initial bases of the second exon but is 5 bases from the 5' end.

### **Detailed description of the invention**

The primary functional role of IgA antibodies is to protect epithelial surfaces from infectious agents. Therefore, the principal sites of IgA synthesis and secretion are the gut, the respiratory epithelium, the lactating breast and other exocrine glands such as the salivary and tear glands. IgA antibodies are selectively transported across epithelia into sites such as the lumen of the gut, where they neutralise toxins and viruses and block the entry of bacteria across the intestinal epithelium. IgA antibodies are also secreted in breast milk (colostrum) and transferred to the gut of newborn offspring to provide protection against disease.

The IgA molecule is composed of two light chains and two heavy ( $\alpha$ ) chains. The heavy chain is divided into one variable domain ( $V_H$ ) and three constant domains ( $C_H1$ ,  $C_H2$  and  $C_H3$ ). The heavy chain also comprises a flexible stretch of polypeptide chain known as the hinge region, which is located between the  $C_H1$  and  $C_H2$  domains. The flexibility of the hinge region is required to allow both arms of the antibody

molecule to bind sites that are different distances apart, and is also required for interaction with antibody-binding proteins that mediate immune effector mechanisms.

The present inventors have identified four novel allelic variants (allotypes) of canine IgA. These variants differ in the sequence of the  $\alpha$  heavy chain gene, and in particular, in the coding region of the hinge. These differences result in variation in hinge length between each allelic variant. This variation is in part due to the presence of a single nucleotide polymorphism in the splice acceptor for the second exon at position 547 in relation to SEQ ID NO: 1. The presence of deoxythreonine (T) at position 547 results in a short hinge variant, such as variants A and B. The presence of deoxyadenosine (A) at position 547 abolishes the splice acceptor site, and therefore results in long hinge variants such as variants C and D. Variation in hinge length is also caused by other sequence differences in the coding region of the hinge, as discussed herein.

The presence of an extended hinge region of IgA makes it more susceptible to cleavage by proteases. Various pathogens secrete proteases that can cleave the hinge region of Ig molecules. Therefore, the presence of a long hinge variant of IgA in an animal may cause susceptibility to disease caused by such pathogens, for example gastrointestinal, skin and respiratory disease.

Accordingly, the present invention provides a method for determining susceptibility to an IgA-related disorder in an animal, the method comprising:

- a) identifying the or each IgA allelic variant present in an animal; and
- b) thereby determining whether the animal is susceptible to an IgA-related disorder.

The IgA-related disorder may be any disease, condition or disorder that is associated with IgA deficiency or dysfunction. Such a disorder is typically an immune related disorder or a bacterial overgrowth disorder in any epithelium where IgA is secreted, such as in the gut, skin or respiratory epithelium. Therefore, the IgA-related disorder is typically a gastrointestinal, skin, respiratory, rheumatoid or periodontal disease. In particular, the disease may be diarrhoea, small intestinal bacterial overgrowth, inflammatory bowel disease, perianal fistulas, atopic dermatitis, pyoderma, anal furunculosis, malassezia infestans or disseminated aspergillosis.

The animal tested is typically a mammal, preferably a non-human animal, such as a dog, cat, horse, pig, cattle or sheep. The animal may be a companion animal or pet. In a preferred embodiment, the animal tested is a dog. The dog tested may be of any

breed, or may be a mixed or crossbred dog, or an outbred dog (mongrel). The dog may be of a breed which is prone to IgA deficiency, such as the German Shepherd Dog, beagle, cocker spaniel, Irish Wolfhound, rottweiler or shar pei. Alternatively the dog tested may be of a breed which is particularly susceptible to skin disease, such as the shar pei, west highland white terrier, Labrador retriever, German Shepherd Dog or golden retriever, or a breed which is known to be susceptible to periodontal disease, such as maltese terriers, shih tsu, Yorkshire terriers, poodles and other small breed dogs.

In one embodiment, the dog tested is of a breed that is susceptible to gastrointestinal disease, such as the boxer, standard poodle, Labrador retriever, Golden retriever or Irish Wolfhound. German Shepherd dogs (also known as Alsatians) are particularly susceptible to a number of gastrointestinal diseases. Therefore, in a preferred embodiment, the dog tested by a method of the invention is of the German Shepherd Dog (GSD) breed. In another aspect, the dog may be a crossbred or outbred dog which is the result of a combination of the German Shepherd Dog breed and one or more other breeds.

In one embodiment of the invention, identification of the IgA allelic variant comprises detecting one or more polymorphisms in the hinge region of the IgA allelic variant, or a polymorphism which is in linkage disequilibrium with such a polymorphism. Preferably, such a polymorphism is at any one of the following positions in relation to SEQ ID NO: 1:

- position 179 [C/T];
- position 370 [T/C];
- position 371 [T/C];
- position 372 [C/G];
- position 375 [G/T];
- positions 514 to 546 [number of CT repeats];
- position 547 [T/A];
- position 563 [A/T];
- positions 563 to 571 [deletion];
- positions 576 to 578 [addition];
- position 582 [C/T];
- position 583 [A/G];
- position 584 [T/A];
- position 592 [G/A]; or

- position 606 [G/A];

The present inventors have identified an association between non-specific dietary sensitivity and the presence of one or more variant C alleles. Therefore, in another preferred embodiment, the animal is tested for the presence of one or more C variant alleles, wherein the presence of at least one variant C allele indicates susceptibility to an IgA-related disorder.

The animal is preferably tested before any symptoms of an IgA-related disorder may be detected. The test may therefore be used to detect susceptibility to disease in an animal, in order to allow prevent the development or onset of an IgA-related disorder. Such preventative action may be related to medical treatment, dietary intervention or any other means of preventing or treating an IgA-related disorder as discussed herein. However, the test may also be used to aid or confirm a diagnosis of an IgA-related disorder in an animal.

#### 15 *Detection of allelic variants*

The detection of allelic variants according to the invention may comprise contacting an IgA polynucleotide or protein of the animal with a specific binding agent for an IgA variant and determining whether the agent binds to the polynucleotide or protein, wherein binding of the agent indicates the presence of the IgA variant, and lack of binding of the agent indicates the absence of the IgA variant.

The method is generally carried out *in vitro* on a sample from the animal. The sample typically comprises a body fluid and/or cells of the individual and may, for example, be obtained using a swab, such as a mouth swab. The sample may be a blood, urine, saliva, skin, cheek cell or hair root sample. The sample is typically processed before the method is carried out, for example DNA extraction may be carried out. The polynucleotide or protein in the sample may be cleaved either physically or chemically, for example using a suitable enzyme. In one embodiment the part of polynucleotide in the sample is copied or amplified, for example by cloning or using a PCR based method prior to detecting the allelic variant(s).

In the present invention, any one or more methods may comprise determining the presence or absence of one or more IgA variants in the animal. The IgA variant is typically detected by directly determining the presence of the polymorphic sequence in a polynucleotide or protein of the animal. Such a polynucleotide is typically genomic

DNA, mRNA or cDNA. The allelic variant may be detected by any suitable method such as those mentioned below.

A specific binding agent is an agent that binds with preferential or high affinity to the protein or polypeptide having the allelic variant but does not bind or binds with only low affinity to other polypeptides or proteins. The specific binding agent may be a probe or primer. The probe may be a protein (such as an antibody) or an oligonucleotide. The probe may be labelled or may be capable of being labelled indirectly. The binding of the probe to the polynucleotide or protein may be used to immobilise either the probe or the polynucleotide or protein.

Generally in the method, determination of the binding of the agent to the IgA variant can be carried out by determining the binding of the agent to the polynucleotide or protein of the animal. However in one embodiment the agent is also able to bind the corresponding wild-type sequence, for example by binding the nucleotides or amino acids which flank the allelic variant position, although the manner of binding to the wild-type sequence will be detectably different to the binding of a polynucleotide or protein containing the allelic variant.

The method may be based on an oligonucleotide ligation assay in which two oligonucleotide probes are used. These probes bind to adjacent areas on the polynucleotide which contains the allelic variant, allowing after binding the two probes to be ligated together by an appropriate ligase enzyme. However the presence of single mismatch within one of the probes may disrupt binding and ligation. Thus ligated probes will only occur with a polynucleotide that contains the allelic variant, and therefore the detection of the ligated product may be used to determine the presence of the allelic variant.

In one embodiment the probe is used in a heteroduplex analysis based system. In such a system when the probe is bound to polynucleotide sequence containing the allelic variant it forms a heteroduplex at the site where the allelic variant occurs and hence does not form a double strand structure. Such a heteroduplex structure can be detected by the use of single or double strand specific enzyme. Typically the probe is an RNA probe, the heteroduplex region is cleaved using RNAase H and the allelic variant is detected by detecting the cleavage products.

The method may be based on fluorescent chemical cleavage mismatch analysis which is described for example in PCR Methods and Applications 3, 268-71 (1994) and Proc. Natl. Acad. Sci. 85, 4397-4401 (1998).

In one embodiment a PCR primer is used that primes a PCR reaction only if it binds a polynucleotide containing the allelic variant, for example a sequence- or allele-specific PCR system, and the presence of the allelic variant may be determined by the detecting the PCR product. Preferably the region of the primer which is complementary to the allelic variant is at or near the 3' end of the primer. The presence of the allelic variant may be determined using a fluorescent dye and quenching agent-based PCR assay such as the Taqman PCR detection system. In a preferred embodiment, one or more of the probes and/or primers shown in Table 4 (for example, all of the probes and primers in Table 4) are used in a Taqman assay to detect an allelic variant.

The specific binding agent may be capable of specifically binding the amino acid sequence encoded by a variant sequence. For example, the agent may be an antibody or antibody fragment. The detection method may be based on an ELISA system. The method may be an RFLP based system. This can be used if the presence of the allelic variant in the polynucleotide creates or destroys a restriction site that is recognised by a restriction enzyme.

The presence of the allelic variant may be determined based on the change which the presence of the allelic variant makes to the mobility of the polynucleotide or protein during gel electrophoresis. In the case of a polynucleotide single-stranded conformation allelic variant (SSCP) or denaturing gradient gel electrophoresis (DDGE) analysis may be used. In another method of detecting the allelic variant a polynucleotide comprising the polymorphic region is sequenced across the region which contains the allelic variant to determine the presence of the allelic variant.

### *Polynucleotides*

The invention also provides a polynucleotide which comprises an IgA variant sequence. An IgA variant sequence typically differs from SEQ ID NO: 1 (Genbank sequence L36871) at one or more of the following polymorphic positions:

- position 179 [C/T];
- position 370 [T/C];
- position 371 [T/C];
- position 372 [C/G];
- position 375 [G/T];
- positions 514 to 546 [number of CT repeats];
- position 547 [T/A];

- position 563 [A/T];
- positions 563 to 571 [deletion];
- positions 576 to 578 [addition];
- position 582 [C/T];
- 5 - position 583 [A/G];
- position 584 [T/A];
- position 592 [G/A]; or
- position 606 [G/A];

Preferably the allelic variant sequence is variant A, B, C or D as discussed  
 10 herein. Accordingly, a polynucleotide of the invention preferably comprises the  
 sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7 or SEQ ID NO: 9 or a  
 fragment thereof. The polynucleotide is typically at least 10, 15, 20, 30, 50, 100, 200 or  
 500 bases long, such as at least or up to 1kb, 10kb, 100kb, 1000 kb or more in length.  
 The polynucleotide will typically comprise flanking nucleotides on one or both sides of  
 15 (5' or 3' to) the allelic variant, for example at least 2, 5, 10, 15 or more flanking  
 nucleotides in total or on each side. Typically, the polynucleotide will be at least 95%,  
 preferably at least 99%, even more preferably at least 99.9% identical to the  
 polynucleotide sequences of SEQ ID NO: 3, 5, 7 or 9. Such numbers of substitutions  
 and/or insertions and/or deletions and/or percentage identity may be taken over the  
 20 entire length of the polynucleotide or over 50, 30, 15, 10 or less flanking nucleotides in  
 total or on each side.

The polynucleotide may be RNA or DNA, including genomic DNA, synthetic  
 DNA or cDNA. The polynucleotide may be single or double stranded. The  
 polynucleotide may comprise synthetic or modified nucleotides, such as  
 25 methylphosphonate and phosphorothioate backbones or the addition of acridine or  
 polylysine chains at the 3' and/or 5' ends of the molecule.

A polynucleotide of the invention may be used as a primer, for example for  
 PCR, or a probe. A polynucleotide or polypeptide of the invention may carry a  
 revealing label. Suitable labels include radioisotopes such as <sup>32</sup>P or <sup>35</sup>S, fluorescent  
 30 labels, enzyme labels or other protein labels such as biotin.

The invention also provides expression vectors that comprise polynucleotides of  
 the invention and are capable of expressing a polypeptide of the invention. Such  
 vectors may also comprise appropriate initiators, promoters, enhancers and other  
 elements, such as for example polyadenylation signals which may be necessary, and

which are positioned in the correct orientation, in order to allow for protein expression. Thus the coding sequence in the vector is operably linked to such elements so that they provide for expression of the coding sequence (typically in a cell). The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner.

The vector may be for example plasmid, virus or phage vector. Typically the vector has an origin of replication. The vector may comprise one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a resistance gene for a fungal vector. Vectors may be used *in vitro*, for example for the production of DNA or RNA or used to transfect or transform a host cell, for example, a mammalian host cell. The vectors may also be adapted to be used *in vivo*, for example in a method of gene therapy.

Promoters and other expression regulation signals may be selected to be compatible with the host cell for which expression is designed. For example, yeast promoters include *S. cerevisiae* GAL4 and ADH promoters, *S. pombe* *nmt1* and *adh* promoter. Mammalian promoters include the metallothionein promoter which can be induced in response to heavy metals such as cadmium. Viral promoters such as the SV40 large T antigen promoter or adenovirus promoters may also be used. Mammalian promoters, such as  $\beta$ -actin promoters, may be used. Tissue-specific promoters are especially preferred. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR), the rous sarcoma virus (RSV) LTR promoter, the SV40 promoter, the human cytomegalovirus (CMV) IE promoter, adenovirus, HSV promoters (such as the HSV IE promoters), or HPV promoters, particularly the HPV upstream regulatory region (URR).

The vector may further include sequences flanking the polynucleotide giving rise to polynucleotides which comprise sequences homologous to eukaryotic genomic sequences, preferably mammalian genomic sequences, or viral genomic sequences. This will allow the introduction of the polynucleotides of the invention into the genome of eukaryotic cells or viruses by homologous recombination. In particular, a plasmid vector comprising the expression cassette flanked by viral sequences can be used to prepare a viral vector suitable for delivering the polynucleotides of the invention to a mammalian cell. Other examples of suitable viral vectors include herpes simplex viral vectors and retroviruses, including lentiviruses, adenoviruses, adeno-associated viruses and HPV viruses. Gene transfer techniques using these viruses are known to those



skilled in the art. Retrovirus vectors for example may be used to stably integrate the polynucleotide giving rise to the polynucleotide into the host genome. Replication-defective adenovirus vectors by contrast remain episomal and therefore allow transient expression.

Polynucleotides of the invention may be used as a probe or primer which is capable of selectively binding to an IgA variant. Preferably the probe or primer is capable of selectively binding to the polynucleotide sequence of SEQ ID NO: 3, 5, 7 or 9. The invention thus provides a probe or primer for use in a method according to the invention, which probe or primer is capable of selectively detecting the presence of an IgA variant. Preferably the probe is isolated or recombinant nucleic acid. It may correspond to or be antisense to the polynucleotide sequence of SEQ ID NO: 3, 5, 7 or 9. The probe may be immobilised on an array, such as a polynucleotide array.

Such primers, probes and other fragments will preferably be at least 10, preferably at least 15 or at least 20, for example at least 25, at least 30 or at least 40 nucleotides in length. They will typically be up to 40, 50, 60, 70, 100 or 150 nucleotides in length. Probes and fragments can be longer than 150 nucleotides in length, for example up to 200, 300, 400, 500, 600, 700 nucleotides in length, or even up to a few nucleotides, such as five or ten nucleotides, short of a full length polynucleotide sequence of the invention.

### Homologues

Homologues of polynucleotide or protein sequences are referred to herein. Such homologues typically have at least 70% homology, preferably at least 80, 90%, 95%, 97% or 99% homology, for example over a region of at least 15, 20, 30, 100 more contiguous nucleotides or amino acids. The homology may be calculated on the basis of nucleotide or amino acid identity (sometimes referred to as "hard homology").

For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (such as identifying equivalent or corresponding sequences (typically on their default settings), for example as described in Altschul S. F. (1993) *J Mol Evol* 36:290-300; Altschul, S, F *et al* (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length  $W$  in the query sequence that either match or satisfy some positive-valued threshold score  $T$  when aligned with a word of the same length in a database sequence.  $T$  is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity  $X$  from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters  $W$ ,  $T$  and  $X$  determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length ( $W$ ) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments ( $B$ ) of 50, expectation ( $E$ ) of 10,  $M=5$ ,  $N=4$ , and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability ( $P(N)$ ), which provides an indication of the probability by which a match between two polynucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The homologous sequence typically differs by at least 1, 2, 5, 10, 20 or more mutations, which may be substitutions, deletions or insertions of nucleotide or amino acids. These mutations may be measured across any of the regions mentioned above in relation to calculating homology. In the case of proteins the substitutions are preferably conservative substitutions. These are defined according to the following Table. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar – uncharged	C S T M
		N Q
	Polar – charged	D E
		K R
AROMATIC		H F W Y

Shorter polypeptide sequences are also within the scope of the invention. For example, a fragment of a polypeptide sequence of the invention is typically at least 10, 15, 20, 30, 40, 50, 60, 70, 80, 100, 150 or 200 amino acids in length. In particular, this aspect of the invention encompasses the situation where the polypeptide is a fragment of a variant canine IgA heavy  $\alpha$  chain which differs in amino acid sequence to a corresponding fragment of the non-variant sequence (i.e. SEQ ID NO: 1). A fragment of the variant IgA may be a Fv, F(ab') or F(ab')<sub>2</sub> fragment. A variant IgA fragment of the invention typically comprises the hinge region of the heavy chain.

Polypeptides of the invention may be chemically modified, for example post-translationally modified. The polypeptides may be glycosylated or comprise modified amino acid residues. Such modified polypeptides fall within the scope of the term "polypeptide" of the invention.

The polypeptides, polynucleotides, vectors, cells or antibodies of the invention may be present in an isolated or substantially purified form. They may be mixed with carriers or diluents which will not interfere with their intended use and still be regarded as substantially isolated. They may also be in a substantially purified form, in which case they will generally comprise at least 90%, e.g. at least 95%, 98% or 99%, of the proteins, polynucleotides, cells or dry mass of the preparation.

It is understood that any of the above features that relate to polynucleotides and proteins may also be a feature of the other polypeptides and proteins mentioned herein, such as the polypeptides and proteins used in the screening and therapeutic aspects of the invention. In particular such features may be any of the lengths, modifications and vectors forms mentioned above.

### *Detector antibodies*

The invention also provides detector antibodies that are specific for a polypeptide of the invention. A detector antibody is specific for one IgA variant, for example, variant A, B, C or D, but does not bind to any other IgA variant. The detector antibodies of the invention are for example useful in purification, isolation or screening methods involving immunoprecipitation techniques.

Antibodies may be raised against specific epitopes of the polypeptides of the invention. An antibody, or other compound, "specifically binds" to a polypeptide when it binds with preferential or high affinity to the protein for which it is specific but does not bind or binds with only low affinity to other polypeptides. A variety of protocols for competitive binding or immunoradiometric assays to determine the specific binding capability of an antibody are well known in the art (see for example Maddox *et al*, J. Exp. Med. 158, 1211-1226, 1993). Such immunoassays typically involve the formation of complexes between the specific protein and its antibody and the measurement of complex formation.

For the purposes of this invention, the term "antibody", unless specified to the contrary, includes fragments which bind a polypeptide of the invention. Such fragments include Fv, F(ab') and F(ab')<sub>2</sub> fragments, as well as single chain antibodies. Furthermore, the antibodies and fragment thereof may be chimeric antibodies, CDR-grafted antibodies or humanised antibodies.

Antibodies may be used in a method for detecting polypeptides of the invention in a biological sample (such as any such sample mentioned herein), which method comprises:

- I providing an antibody of the invention;
- II incubating a biological sample with said antibody under conditions which allow for the formation of an antibody-antigen complex; and
- III determining whether antibody-antigen complex comprising said antibody is formed.

Antibodies of the invention can be produced by any suitable method. Means for preparing and characterising antibodies are well known in the art, see for example Harlow and Lane (1988) "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. For example, an antibody may be produced by raising antibody in a host animal against the whole polypeptide or a fragment thereof, for example an antigenic epitope thereof, herein after the "immunogen". The

fragment may be any of the fragments mentioned herein (typically at least 10 or at least 15 amino acids long).

A method for producing a polyclonal antibody comprises immunising a suitable host animal, for example an experimental animal, with the immunogen and isolating immunoglobulins from the animal's serum. The animal may therefore be inoculated with the immunogen, blood subsequently removed from the animal and the IgG fraction purified. A method for producing a monoclonal antibody comprises immortalising cells which produce the desired antibody. Hybridoma cells may be produced by fusing spleen cells from an inoculated experimental animal with tumour cells (Kohler and Milstein (1975) *Nature* 256, 495-497).

An immortalized cell producing the desired antibody may be selected by a conventional procedure. The hybridomas may be grown in culture or injected intraperitoneally for formation of ascites fluid or into the blood stream of an allogenic host or immunocompromised host. Human antibody may be prepared by *in vitro* immunisation of human lymphocytes, followed by transformation of the lymphocytes with Epstein-Barr virus.

For the production of both monoclonal and polyclonal antibodies, the experimental animal is suitably a goat, rabbit, rat, mouse, guinea pig, chicken, sheep or horse. If desired, the immunogen may be administered as a conjugate in which the immunogen is coupled, for example via a side chain of one of the amino acid residues, to a suitable carrier. The carrier molecule is typically a physiologically acceptable carrier. The antibody obtained may be isolated and, if desired, purified.

#### *Detection kit*

The invention also provides a kit that comprises means for determining the presence or absence of one or more IgA allelic variant(s) in an animal. In particular, such means may include a specific binding agent, probe, primer, pair or combination of primers, or antibody, including an antibody fragment, as defined herein which is capable of detecting or aiding detection of an IgA allelic variant. The primer or pair or combination of primers may be sequence specific primers which only cause PCR amplification of a polynucleotide sequence comprising the IgA variant to be detected, as discussed herein. The kit may also comprise a specific binding agent, probe, primer, pair or combination of primers, or antibody which is capable of detecting the absence of the allelic variant. The kit may further comprise buffers or aqueous solutions.

The kit may additionally comprise one or more other reagents or instruments which enable any of the embodiments of the method mentioned above to be carried out. Such reagents or instruments may include one or more of the following: a means to detect the binding of the agent to the allelic variant, a detectable label such as a  
 5 fluorescent label, an enzyme able to act on a polynucleotide, typically a polymerase, restriction enzyme, ligase, RNase H or an enzyme which can attach a label to a polynucleotide, suitable buffer(s) or aqueous solutions for enzyme reagents, PCR primers which bind to regions flanking the allelic variant as discussed herein, a positive and/or negative control, a gel electrophoresis apparatus, a means to isolate DNA from  
 10 sample, a means to obtain a sample from the individual, such as swab or an instrument comprising a needle, or a support comprising wells on which detection reactions can be carried out. The kit may be, or include, an array such as a polynucleotide array comprising the specific binding agent, preferably a probe, of the invention. The kit typically includes a set of instructions for using the kit.

#### Screening for therapeutic agents

The present invention also relates to the use of variant IgA as a screening target for identifying therapeutic agents for the treatment of IgA-related disorders. In one embodiment the invention provides a method for identifying an agent useful for the  
 20 treatment of IgA-related disorders, which method comprises contacting a variant IgA polypeptide or a polynucleotide with a test agent and determining whether the agent is capable of binding to the polypeptide or modulating the activity or expression of the polypeptide or polynucleotide. Any suitable binding assay format can be used to determine whether the IgA variant binds the test agent, such as the formats discussed  
 25 below.

The method may be carried out *in vitro*, either inside or outside a cell, or *in vivo*. In one embodiment the method is carried out on a cell, cell culture or cell extract that comprises a variant IgA protein or polynucleotide. The cell may be any suitable cell, and is typically a cell in which the product is naturally expressed. For example, the cell  
 30 may be a mucosal epithelial cell such as an IgA<sup>+</sup> plasma cell from the duodenal lamina propria. The method may also be carried out *in vivo* in a non-human animal which is transgenic for an IgA variant polynucleotide. The transgenic non-human animal is typically of a species commonly used in biomedical research and is preferably a

laboratory strain. Suitable animals include rodents, particularly a mouse, rat, guinea pig, ferret, gerbil or hamster. Most preferably the animal is a mouse.

The term "modulate" includes any of the ways mentioned herein in which the agent is able to modulate activity of an IgA variant polypeptide or polynucleotide. This may be determined by contacting the polypeptide or polynucleotide with the test agent under conditions that permit activity of the polypeptide or polynucleotide, and determining whether the test agent is able to modulate the activity of the polypeptide or polynucleotide.

The activity which is measured may be any of the activities which are mentioned herein, and may be the measurement of a property of the IgA variant polypeptide or polynucleotide, or an effect on a cellular component, cell or animal in which the method is being carried out. The effect may be one that is associated with an IgA-related disorder, and may be a characteristic or symptom of an IgA-related disorder, such as any such characteristic or symptom mentioned herein.

In one embodiment the assay measures the effect of the test agent on the binding between the variant IgA polypeptide or polynucleotide and another agent, such as a protease. In particular, the assay may include proteases from pathogens that are known to cause IgA-related disorders in dogs. Suitable assays in order to measure the changes in such interactions include fluorescence imaging plate reader assays, and radioligand binding assays. In the case where the activity is transcription from a gene the method may comprise measuring the ability of the candidate substance to modulate transcription, for example in a reporter gene assay.

Suitable candidate agents which may be tested in the above screening methods include antibody agents, for example monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies and CDR-grafted antibodies. Furthermore, combinatorial libraries, defined chemical identities, peptide and peptide mimetics, oligonucleotides and natural agent libraries, such as display libraries may also be tested. The test agents may be chemical compounds, which are typically derived from synthesis around small molecules which may have any of the properties of the agent mentioned herein. Batches of the candidate agents may be used in an initial screen of, for example, ten substances per reaction, and the substances of batches which show modulation tested individually. The term 'agent' is intended to include a single substance and a combination of two, three or more substances. For example, the term agent may refer to

a single peptide, a mixture of two or more peptides or a mixture of a peptide and a defined chemical entity.

In one aspect of the invention, the test agent is a food ingredient. Hence, the invention relates to a method of screening food ingredients to determine whether they contribute to or aggravate gastrointestinal disease in susceptible dogs, or if they prevent or alleviate gastrointestinal disease. The food ingredient may be one that is typically used in dog or other types of food, or may be a novel food ingredient.

The present invention also provides an agent identified by a screening method of the invention. An agent identified in the screening method of the invention may be used in the therapeutic treatment of an IgA-related disorder. Such an agent may be formulated and administered in any means or amounts as discussed below.

#### *Treatment of IgA-related disorders*

The invention provides a method of treating an animal for an IgA-related disorder, the method comprising identifying an animal which is susceptible to an IgA-related disorder by a method of the invention, and administering to the animal an effective amount of a therapeutic agent which treats the IgA-related disorder. The IgA-related disorder may be any disease or disorder mentioned herein, and is typically a gastrointestinal, skin, respiratory disease, rheumatoid or periodontal disease. The therapeutic agent is typically a drug such as an anti-inflammatory (e.g. sulphur salazine), a corticosteroid (e.g. prednisolone), an antibiotic (e.g. amoxycillin or enrofloxacin) or a protease inhibitor (e.g. amprenavir). The therapeutic agent may be any drug known in the art that may be used to treat an IgA-related disorder, or may an agent identified by a screening method as discussed previously.

The therapeutic treatment may result in a change of the bacterial flora of the animal. In particular, the bacterial flora of the animal may be altered to reduce the production of proteases that can degrade IgA. Such a change is typically effected in the gastrointestinal system of the animal, and may be carried out by administering agents such as prebiotics or probiotics to the animal.

The therapeutic agent may be administered in various manners such as orally, intracranially, intravenously, intramuscularly, intraperitoneally, intranasally, intrademally, and subcutaneously. The pharmaceutical compositions that contain the therapeutic agent will normally be formulated with an appropriate pharmaceutically acceptable carrier or diluent depending upon the particular mode of administration



being used. For instance, parenteral formulations are usually injectable fluids that use pharmaceutically and physiologically acceptable fluids such as physiological saline, balanced salt solutions, or the like as a vehicle. Oral formulations, on the other hand, may be solids, for example tablets or capsules, or liquid solutions or suspensions. In a preferred embodiment, the therapeutic agent is administered to the animal in its diet, for example in its drinking water or food.

The amount of therapeutic agent that is given to an animal will depend upon a variety of factors including the condition being treated, the nature of the animal under treatment and the severity of the condition under treatment. A typical daily dose is from about 0.1 to 50 mg per kg, preferably from about 0.1mg/kg to 10mg/kg of body weight, according to the activity of the specific inhibitor, the age, weight and conditions of the animal to be treated, the type and severity of the disease and the frequency and route of administration. Preferably, daily dosage levels are from 5 mg to 2 g.

#### *Customised food*

In one aspect, the invention relates to a customised diet for an animal that is susceptible to an IgA-related disorder. In a preferred embodiment, the customised food is for a companion animal or pet, such as a dog. Such a food may be in the form of, for example, wet pet foods, semi-moist pet foods, dry pet foods and pet treats. Wet pet food generally has a moisture content above 65%. Semi-moist pet food typically has a moisture content between 20-65% and can include humectants and other ingredients to prevent microbial growth. Dry pet food, also called kibble, generally has a moisture content below 20% and its processing typically includes extruding, drying and/or baking in heat. The ingredients of a dry pet food generally include cereal, grains, meats, poultry, fats, vitamins and minerals. The ingredients are typically mixed and put through an extruder/cooker. The product is then typically shaped and dried, and after drying, flavours and fats may be coated or sprayed onto the dry product.

Accordingly, the present invention enables the preparation of customised food suitable for an animal which is susceptible to an IgA-related disorder, wherein the customised animal food formulation comprises ingredients that prevent or alleviate IgA-related disorders, and/or does not comprise components that contribute to or aggravate IgA-related disorders. Such ingredients may be any of those known in the art to prevent or alleviate an IgA-related disorder. Alternatively, such ingredients may be identified

by screening methods as discussed herein. The preparation of customised animal food may be carried out by electronic means, for example by using a computer system.

In one embodiment, the customised food may be formulated to alter the profile of food proteins in order to minimise the potential for secondary dietary sensitivity.

5 The customised food may be hypoallergenic or may exclude ingredients that are poorly tolerated or cause allergies, for example gluten-containing grains such as wheat, particular protein sources such as animal proteins, milk (lactose), eggs, soy, peanuts, shellfish, fruits or tree nuts.

In another embodiment, the customised food may be formulated to include  
10 functional ingredients that help prevent or alleviate an IgA-related disorder. Such a functional ingredient may be a compound that stimulates immune function or protects against degradation of IgA, for example  $\beta$ -glucans or glutamine. Alternatively, exogenous IgA may be administered orally in the diet, for example using colostrum or eggs. These may be hyperimmunised to pathogens, such as pathogenic *E. coli* (e.g.  
15 EPEC), *Campylobacter* or *Salmonella*.

The functional ingredient may help to prevent or alleviate an IgA-related disorder by improving gut barrier function, for example prebiotics, probiotics or oligosaccharides. In one aspect, the customised food is formulated to prevent or  
20 alleviate IgA-related skin disease. Such food may comprise functional ingredients that improve the condition of the skin, for example vitamin C, taurine, curcumin or aloe vera. One particular example of a skin support diet is described in International Patent Application No. PCT/GB02/02538.

The present invention also relates to a method of providing a customised animal food, comprising providing food suitable for an animal which is susceptible to an IgA-  
25 related disorder to the animal, the animal's owner or the person responsible for feeding the animal, wherein the animal has been determined to be susceptible to an IgA-related disorder by a method of the invention.

### *Bioinformatics*

30 The sequences of the IgA variants may be stored in an electronic format, for example in a computer database. Accordingly, the invention provides a database comprising information relating to IgA allelic variant sequences. The database may include further information about the allelic variant, for example the level of association of the allelic variant with an IgA-related disorder or the frequency of the allelic variant

in the population. In one aspect of the invention, the database further comprises information regarding the food components which are suitable and the food components which are not suitable for animals who possess a particular allelic variant of IgA.

A database as described herein may be used to determine the susceptibility of an animal to an IgA-related disorder. Such a determination may be carried out by electronic means, for example by using a computer system (such as a PC). Typically, the determination will be carried out by inputting genetic data from the animal to a computer system; comparing the genetic data to a database comprising information relating to IgA allelic variants; and on the basis of this comparison, determining the susceptibility of the animal to an IgA-related disorder.

The invention also provides a computer program comprising program code means for performing all the steps of a method of the invention when said program is run on a computer. Also provided is a computer program product comprising program code means stored on a computer readable medium for performing a method of the invention when said program is run on a computer. A computer program product comprising program code means on a carrier wave that, when executed on a computer system, instruct the computer system to perform a method of the invention is additionally provided.

As illustrated in Figure 6, the invention also provides an apparatus arranged to perform a method according to the invention. The apparatus typically comprises a computer system, such as a PC. In one embodiment, the computer system comprises: means 20 for receiving genetic data from the animal; a module 30 for comparing the data with a database 10 comprising information relating to IgA allelic variants; and means 40 for determining on the basis of said comparison the susceptibility of the animal to an IgA-related disorder.

#### *Food manufacturing*

In one embodiment of the invention, the manufacture of a customised animal food may be controlled electronically. Typically, information relating to the IgA allelic variant(s) present in an animal may be processed electronically to generate a customised animal food formulation. The customised animal food formulation may then be used to generate electronic manufacturing instructions to control the operation of food manufacturing apparatus. The apparatus used to carry out these steps will typically comprise a computer system, such as a PC, which comprises means 50 for processing

the nutritional information to generate a customised animal food formulation; means 60 for generating electronic manufacturing instructions to control the operation of food manufacturing apparatus; and a food product manufacturing apparatus 70.

5 The food product manufacturing apparatus used in the present invention typically comprises one or more of the following components: container for dry pet food ingredients; container for liquids; mixer; former and/or extruder; cut-off device; cooking means (e.g. oven); cooler; packaging means; and labelling means. A dry ingredient container typically has an opening at the bottom. This opening may be covered by a volume-regulating element, such as a rotary lock. The volume-regulating  
10 element may be opened and closed according to the electronic manufacturing instructions to regulate the addition of dry ingredients to the pet food.

Dry ingredients typically used in the manufacture of pet food include corn, wheat, meat and/or poultry meal. Liquid ingredients typically used in the manufacture of pet food include fat, tallow and water. A liquid container may contain a pump that  
15 can be controlled, for example by the electronic manufacturing instructions, to add a measured amount of liquid to the pet food.

In one embodiment, the dry ingredient container(s) and the liquid container(s) are coupled to a mixer and deliver the specified amounts of dry ingredients and liquids to the mixer. The mixer may be controlled by the electronic manufacturing instructions.  
20 For example, the duration or speed of mixing may be controlled. The mixed ingredients are typically then delivered to a former or extruder. The former/extruder may be any former or extruder known in the art that can be used to shape the mixed ingredients into the required shape. Typically, the mixed ingredients are forced through a restricted opening under pressure to form a continuous strand. As the strand is extruded, it may  
25 be cut into pieces (kibbles) by a cut-off device, such as a knife. The kibbles are typically cooked, for example in an oven. The cooking time and temperature may be controlled by the electronic manufacturing instructions. The cooking time may be altered in order to produce the desired moisture content for the food. The cooked kibbles may then be transferred to a cooler, for example a chamber containing one or  
30 more fans.

The food manufacturing apparatus may comprise a packaging apparatus. The packaging apparatus typically packages the food into a container such as a plastic or paper bag or box. The apparatus may also comprise means for labelling the food, typically after the food has been packaged. The label may provide information such as:

ingredient list; nutritional information; date of manufacture; best before date; weight; and species and/or breed(s) for which the food is suitable.

The invention is illustrated by the following Examples:

## 5 Example 1

### **Materials and Methods**

#### *Sample Collection*

Endoscopic biopsies of duodenal mucosa were obtained from dogs presented to  
10 the Department of Clinical Veterinary Science, University of Bristol for investigation of gastrointestinal disease. Gastroduodenoscopy was performed using a GIF-XQ230 flexible video endoscope (Olympus Keymed, Southend-on-Sea, UK). Multiple mucosal biopsies were taken at the level of the caudal duodenal flexure using FB-25K biopsy forceps (Olympus Keymed). Biopsies were placed in a 1.0-ml cryotube (NUNC,  
15 Fischer Scientific Ltd., Loughborough, Leicestershire), snap frozen in liquid nitrogen and stored at -70°C.

#### *RNA/DNA Isolation*

Two endoscopic biopsies (total tissue mass 9-16mg) were added to a green  
20 Ribolyser tube (Ribolyser System, Thermo-Hybaid, Ashford, Middlesex, UK) containing 400µl of lysis buffer from the RNA isolation kit (see below) and processed for 45 seconds at 6.0 m/s to homogenise the biopsies. This lysate was processed through the RNeasy Isolation System (Qiagen Ltd., Crawley, U.K.) as per the manufacturer's protocol except that the RNA was eluted in 100µl of nuclease-free  
25 waster. This procedure produces a mixture of total RNA and a significant amount of genomic DNA. A negative control of nuclease free water was passed through the extraction procedure. Samples were stored at -70°C prior to use.

#### *DNase Digestion*

30 DNase digestion was carried out on 50µl of the RNA/DNA mix from the extraction detailed above. DNase digestion was carried out both in solution and on the subsequent Qiagen column used to further purify the RNA. In-solution DNase digestion was carried out by treating 50µl of the extraction mix with 5units of

Amplification Grade DNase 1 (Invitrogen Ltd.) as per the manufacturer's instructions. In order to remove any residual DNase or EDTA from the treated RNA, the solution was passed through the RNeasy Isolation System (Qiagen Ltd.) a second time using the RNA clean-up protocol. A second DNase digestion was carried out on this column using the RNase-Free DNase Set (Qiagen Ltd.).

### Primer Design

Primers for production of the cloned RT-PCR and PCR products were designed using Primer 3 ([www-genome.wi.mit.edu/cgi-bin/primer/primer3\\_www.cgi](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi)) against the Genbank sequence for canine  $\alpha$ -chain (accession number: L36871) and are shown in Table 1. The forward primer was located in the 3' end of exon 1 and the reverse primer in the 5' end of exon 2 which together produced a 550bp DNA product and a 300bp product with cDNA. The primer and probe sequences for identification of suitable dogs to clone DNA and RNA sequences for  $\alpha$ -chain from have been used previously for the quantification of  $\alpha$ -chain transcript from canine endoscopic duodenal mucosal biopsies and are shown in Table 1 ( $\alpha$ -chain set 1).

Table 1: Primer and Probe Sequences

Primer Set	Product Length	Forward Primer (5'-3')	Reverse Primer (5'-3')	5'-Fluorophore	Probe Sequence (5'-3')	3'-Quencher
Cloned Product Primers	550bp gDNA 300bp cDNA	TGTGAAC GTGACCT GGAATG	GCATTGG AGCCTAA AAGCAG	Texas Red	GTCATCCATG TCCCTCGTGC AATGAG	BHQ-2
$\alpha$ -Chain Set 1	84bp cDNA	TGTGCCCT GCAAAGA TAACA	AGGGCTG GCTTCTGT AGTGA			
$\alpha$ -Chain Set 2	136bp cDNA	CGTCTGTG AAATGCC AAGTG	AGGGCTG GCTTCTGT AGTGA			

This original set of primers was designed such that the forward primer spanned the junction between the first and second exon. A second set of primers was designed using Primer3 with the same probe sequence, such that the forward primer was located in the first exon ( $\alpha$ -chain set 2). The primer and probe set was designed such that the annealing temperatures of the primers were 60°C and the probe 8-10°C higher, and that a product of between 80 and 200 bases in length would be obtained. In order to minimise primer-dimer formation, the maximum self-complementarity was 6 and the maximum 3' self-complementarity was 2.

## 10 *RT-PCR*

Gene specific real-time RT-PCR amplification of  $\alpha$ -chain was performed using the Platinum Quantitative RT-PCR Thermoscript™ One-Step System (Invitrogen Ltd) using 5µl of RNA, 4.5mM Mg<sup>2+</sup> and the primers and probe at a concentration of 200nM and 100nM respectively, in a final volume of 25µl. No-RT reactions were made by substituting the Thermoscript enzyme mix with 2 units of Platinum Taq DNA Polymerase (Invitrogen Ltd.). The negative control from the extraction procedure, as well as a nuclease-free water control, were included with all sample runs. The RT-PCR was performed in an iCycler IQ (Bio-Rad Laboratories Ltd, Hercules, CA) with an initial incubation of 55°C for 20 minutes ( $\alpha$ -chain), followed by 95°C for 5 minutes and then 45 cycles of 95°C for 10 seconds, 60°C for 15 seconds during which the fluorescence data were collected. All reactions were made up on ice and placed in the iCycler held at the initial incubation temperature to minimise primer-dimer formation. The threshold cycle (Ct value) was calculated as the cycle when the fluorescence of the sample exceeded a threshold level corresponding to 10 standard deviations from the mean of the baseline fluorescence.

RT-PCR to produce the products for cloning was performed using the Platinum Quantitative RT-PCR Thermoscript™ One-Step System (Invitrogen Ltd) using 5µl of RNA as described previously. DNA amplification was performed by substituting the Thermoscript enzyme mix with 2 units of Platinum Taq DNA Polymerase (Invitrogen Ltd.) with 5µl of the RNA/DNA mix. The PCR protocol was altered for the longer products so that after the initial incubation, 45 cycles of 95°C for 15 seconds, 60°C for 30 seconds and 72°C for 30 seconds were performed followed by a final incubation of 72°C for 5 minutes. These products were separated using 2% agarose gel

electrophoresis and appropriately sized bands were excised from the gel and then purified using the QIAquick PCR Purification Kit (Qiagen Ltd., Crawley, UK) following the manufacturer's instructions.

## 5 *Cloning and Sequencing of Products*

The purified PCR products were cloned using the TOPO TA Cloning kit (Invitrogen Ltd.) as per the manufacturer's instructions using chemically competent *E. Coli*. Following white/blue colony selection, positive clones were sub-cultured overnight and the plasmids were purified using the Qiagen Plasmid Mini Prep Kit (Qiagen Ltd.) as per the manufacturer's instructions. At least four clones per reaction mix (e.g. DNA and RNA) were purified for each dog and were sent to the Sequencing Service (School of Life Sciences, University of Dundee, Dundee, Scotland) for sequencing. Sequence results were aligned with the Genbank sequence using Omega 2.0 (Accelrys, Cambridge, UK).

15

## **Results**

### *IgA expression*

A bimodal pattern of expression of IgA mRNA from canine duodenal mucosa was previously shown (Figure 1A) using primer set 1 (Peters, I. R., C. R. Helps, R. M. Batt, M. J. Day, and E. J. Hall. 2003. Quantitative real-time RT-PCR measurement of mRNA encoding alpha-chain, pIgR and J-chain from canine duodenal mucosa. *J Immunol Methods* 275:213). To confirm the quantification data, a second set of primers were designed (Table 1). Surprisingly, when primer set 2 was used, the bimodal distribution was lost (Figure 1B). The two primer sets differ only in the position of the forward primer, which spans the junction between exon one and two, as shown in Figure 3. These results indicated that a similar amount of  $\alpha$ -chain mRNA is present in all samples but that the forward primer in set 1 does not detect a significant portion of the mRNA in some samples. Therefore, a primer set was designed which amplified a segment of both gDNA and cDNA which encompassed this region, in order to determine whether there were any sequence differences.



### Sequencing Data

Four alternate sequences for the 5'-end of the second exon of the dog alpha heavy chain gene were identified, three of which have not previously been described (Figure 2). The variants were termed A to D, with variant A being similar to the Genbank sequence and thus the variant that was detected using primer set 1. A single nucleotide polymorphism exists at position 179 which differs between individuals with the same variant but this polymorphism does not alter the encoded amino acid (Figure 4).

A major difference between the variants was the position of the splice acceptor site for the second exon. This difference in the splice acceptor site was due to a single base polymorphism at position 547, that resulted in loss of a splice acceptor site for the second exon (Figure 2). The presence of deoxyadenosine at this point resulted in coding for the mRNA from position 549 (variants C and D), whereas the presence of a deoxythreonine at this position led to coding from position 558 (variants A and B) with a transcript that is nine bases shorter.

The shortest variant was variant B which had the deoxythreonine polymorphism at position 558 but also had a nine base deletion within the 5' end of the second exon (positions 563 to 571) and three additional bases after this deletion (positions 576 to 578). This variant also had other base differences at positions 582, 583, 592 and 606 compared with the other variants, and this resulted in alteration in the predicted amino acid sequence. The major difference between variants C and D was the presence of a three base addition similar to that in variant B between positions 576 to 578, making variant D the longest of the four.

The combinations of variants found in each individual dog are detailed in Table 2. These results indicate that more than one variant can be detected in some individual dogs, suggesting that heterozygous individuals exist. There is also evidence that both variants possessed by an individual are transcribed as the sequences were detected in both gDNA and cDNA products with the exception of \* and +. These were only found in the gDNA (\*) and cDNA (+) products respectively. All variants were found in two or more individuals with the exception of variant D which was only found in dog 48.

Table 2: Variant of IgA found in each dog sequenced

Dog	Breed	Variant A	Variant B	Variant C	Variant D
34	Labrador	-	X	X (*)	-
40	GSD	-	-	X	-
41	GSD	-	-	X	-
43	Staffordshire Bull Terrier	X	X	-	-
45	Greyhound	X	-	X (+)	-
46	Crossbred	X	X	-	-
47	Border Collie	-	-	-	X
48	Cocker Spaniel	-	-	X	

**Example 2**

5

**Materials and Methods***DNA Samples*

Buccal cells were collected from dogs by rotating a sterile cytology brush (Rocket Medical, Cat No. R57483) six times in the inside of the cheek. The brushes were then replaced in their individual wrapper and left to dry for a minimum of two hours at room temperature. DNA was extracted using the Qiagen QIAamp DNA Blood Mini Kit (Cat No. 51104) following the Buccal Swab Spin protocol. The DNA was eluted using 100µl of dH<sub>2</sub>O and then stored at -20°C.

15 *PCR Amplification*

Primers were designed using Primer 3 ([www-genome.wi.mit.edu/cgi-bin/primer/primer3\\_www.cgi](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi)) against the Genbank sequence for the α-chain (accession number: L36871). The forward primer is located in the 3' end of exon 1 and the reverse primer in the 5' end of exon 2 which together produced a 550bp DNA product. The oligos were ordered from Sigma-Genosys, desalted, and used at 0.025 µM synthesis scale. 50µl PCR reactions were carried using 25pmol of each oligo and 50ng of DNA from each dog sample and 25µl of Eurogentec HotGoldstar PCR mastermix containing a red loading dye and 1.5mM MgCl (PK-0073-02R).

Thermal cycling was performed using a Hybrid MBS 0.2S PCR machine using the following cycling conditions: incubation at 95°C for 10 min, followed by 10 cycles of 95°C for 30 sec, 64°C (-1°C per cycle) for 45 sec and 72°C for 90 sec, followed by 28 cycles of 95°C for 30 sec, 55°C for 45 sec and 72°C for 90 sec. 5µl of each of the PCR sample and 1µl of Gelstar nucleic acid gel stain (BioWhittaker Molecular Applications, Cat. No 50535) were run on a 2% agarose (Invitrogen, Cat. No 15510-027) gel at 100mV to check for product. Successful PCR products were purified using a 96 well PCR cleanup plate (Millipore, Cat No MANU03010) following the standard method. The extra wash phase with 50µl of H<sub>2</sub>O was added to remove red dye from the mastermix. Samples were quantified using 1µl of purified PCR product on the Nanodrop Spectrophotometer. Analysis carried out using nucleic acid sample DNA-50 on the Nanodrop 2.4.7a software.

#### *DNA Sequencing*

Cycle sequencing was performed using 25fmol of purified PCR product with the CEQ 2000 Dye Terminator Cycle Sequencing with Quick Start kit (Beckman Coulter, P/N 608120). 20µl reactions were carried out using 3µl of DTCS quick Start Master Mix, 1µl of forward or reverse primer that was used in the PCR step (5pmol) and an adjustable volume of DNA template and dH<sub>2</sub>O. Thermal cycling was performed using the same PCR machine as used in the PCR step but under the following conditions: 30 cycles of 96°C for 20 sec, 50°C for 20 sec and 60°C for 4 min. Following these cycles, the samples were subjected to ethanol precipitation, and were evaporated for dryness using a vacuum pump for 40 min. Samples were resuspended in 40µl of deionised formamide and a drop of mineral oil was placed on top. The samples were run on a Beckman CEQ 2000 Sequencer using the LFR capillary method. The sequence traces were analysed using the CEQ2000XL DNA Analysis System software Version 4.3.9.

#### **Results**

In order to investigate the distribution of IgA alleles within dog breeds, the IgA genotype of 183 dogs from 11 different breeds was determined. The IgA genotypes for each dog breed are shown in Table 3 below. All 54 German Shepherd dogs tested were homozygous for variant C.

Table 3: Breed study results using sequencing method

Breed	Homozygous				Heterozygous					
	AA	BB	CC	DD	AB	AC	AD	BC	BD	CD
German Shepherd			54							
Labrador retriever		8	1		1		1	5		
Shih Tzu	12	1	2							
Rottweiler	15									
Golden retriever		7	4					4		1
Beagle	1	1	2	6						1
Dobermann	5		10							
Yorkshire Terrier	6	1	4			1		1		1
King Charles Cavalier Spaniel		5			5					
West Highland White Terrier	8		3		1					
American Cocker Spaniel	1		4							

**Example 3**

5

**Materials and Methods***Template*

The allelic identities of canine DNA samples were determined using the 5'-3' exonuclease ("TaqMan") assay. Canine genomic DNA was used as the template. The DNA was used either as a direct isolate, or in the case of very dilute samples, was amplified using GenomiPhi™ DNA Amplification Kit (Amersham Biosciences). Assays were performed using DNA at a concentration of 50-100ng/μl.

10

*Reaction*

15

Reactions were carried out using: 12.5μl TaqMan Universal PCR Master Mix, no AmpErase UNG (Applied Biosystems, Warrington, UK); 1μl each oligonucleotide (10pm/μl stock solution) (Sigma Genosys, Cambs, UK); 1μl TaqMan probe (5pm/μl stock solution) (Sigma Genosys, Cambs, UK); 9.5μl nuclease free water; and 1μl template DNA. The probes were labelled 5' with 6FAM and 3' with TAMRA.

Reactions were performed in an ABI 7700 Sequence Detection System using default PCR conditions, with the exception of the annealing temperature, which was raised to 63°C. Primers and probe sequences are shown in Table 4.

5 Table 4: Primer and Probe sequences

Allele	Forward primer	Reverse primer	Probe
A	GAGGGTGCACACTGA CCTGTT	CACGAGGGACATGGA TGAC	CTCTCTCTGCTCCTGAAG ATAACAGTCATCCGT
B	GCACACTGACCTGTT CCAATCTC	GCACACTGACCTGTT CCAATCTC	ATAACTGTCCTCATCTGTG TCCCTCATGCA
C	GAGGGTGCACACTGA CCTGTT	GGGCTGGCTTCTGTA GTGACA	CTCTCTCAGCTCCTGAAG ATAACTGTCATCCGT
D	TCTCTCTCAGCTCCTG AAGATAACTG	GGGCTGGCTTCTGTA GTGACA	CCGTGTCCTCATCCAAGT CCCTCG

### Results

The IgA genotype of 95 dogs from 10 different breeds was determined by the "TaqMan" assay method. The IgA genotypes for each dog breed are shown in Table 5.

- 10 All 10 German Shepherd dogs tested were homozygous for variant C.

Table 5: Breed study results using "TaqMan" assay

Breed	Homozygous				Heterozygous					
	AA	BB	CC	DD	AB	AC	AD	BC	BD	CD
German Shepherd			10							
Labrador retriever		3		1	1		1	3		
Shih Tzu	7	1	1							
Rottweiler	10									
Golden retriever		4	3					2		1
Beagle		1	1	5						1
Dobermann	3		7							
Yorkshire Terrier	4	1	2	1		1		1		
King Charles Cavalier Spaniel		5			5					
West Highland White Terrier	4		4		1					

#### Example 4

The distribution of IgA genotypes in a panel of dietary sensitive dogs was determined. Dietary sensitive dogs are defined as dogs that produce intermittent loose faeces, respond to dietary manipulation and whose dietary sensitivity is not food specific. Genotyping was carried out using the "TaqMan" assay as described in Example 3.

#### Results

The genotypes of the dietary sensitive panel were compared with the genotypes from the "random" population of dogs from Example 3 (Table 6). The most common genotypes within the panel of dietary sensitive dogs were AC (30%) and BC (35%). A binomial test was carried out to see if there was an association between a particular genotype and dietary sensitivity. Table 7 shows the percentage of dogs in each sample with each allele, and the p-value of the difference between the samples (based on the binomial distribution). Within the panel of dietary sensitive dogs, 80% had one or more variant C allele. In comparison, only 39% of the random population of dogs had one or more variant C allele. The association between dietary sensitivity and the presence of variant C is statistically significant ( $p=0.00$ ). These results indicate that the presence of one or more variant C allele correlates with susceptibility to non-specific dietary sensitivity.

Table 6: Comparison of IgA genotypes

Genotype	Random population	Dietary sensitive panel
AA	28 (29%)	2 (10%)
AB	7 (7%)	0 (0%)
AC	1 (1%)	6 (30%)
AD	1 (1%)	0 (0%)
BB	15 (16%)	2 (10%)
BC	6 (6%)	7 (35%)
CC	28 (29%)	2 (10%)
CD	2 (2%)	1 (5%)
DD	7 (7%)	0 (0%)
Total	95	20

Table 7: Frequency of variant alleles

Allelic variant	Random population	Dietary sensitive panel	P-value
A	39%	40%	0.55
B	29%	45%	0.16
C	39%	80%	0.00
D	11%	5%	0.42

## Discussion

5

Multiple IgA subclasses have been identified in humans, primates and lagomorphs, whereas in mice, cattle and dogs only a single subclass has previously been characterised. The two human subclasses (IgA<sub>1</sub> and IgA<sub>2</sub>) are defined by a difference in the length of the hinge region between the CH<sub>1</sub> and CH<sub>2</sub> domains. The single IgA subclass previously identified in dogs has a hinge region with a predicted amino acid sequence similar to the IgA<sub>1</sub> subclass of humans.

10

The presence of the extended hinge region of human IgA<sub>1</sub> confers greater flexibility to the immunoglobulin molecule, facilitating antigen binding, but makes it more susceptible to cleavage by protease compared with IgA<sub>2</sub>. The shortening of the hinge region that is present in human IgA<sub>2</sub> is due to a 39 base pair deletion from the second exon close to the 5' end (Figure 5). This is not caused by a shift in the splice site but is due to a separate gene locus within the immunoglobulin cluster.

15

Of the four variants identified in canine IgA, variants A, C and D share the greatest sequence similarity, with the difference in mRNA length due to a polymorphism in the splice acceptor. Variant B has the shortest mRNA sequence, similar to variant A, but it also has base deletions towards the 5' end of the second exon with a greater number of base polymorphisms compared with the other variants. It is therefore possible that variants A, C and D are allelic variants at one gene locus (similar to human IgA<sub>1</sub>), and that variant B is encoded within a second locus and may therefore represent a second IgA subclass (similar to IgA<sub>2</sub>).

20  
25

German Shepherd dogs (GSD) are particularly prone to a number of inflammatory and immune-mediated alimentary diseases. All sixteen GSD analysed in a separate study had 'low' expression of mRNA when tested with primer set 1, suggesting that none expressed variant A. The susceptibility of GSD to disease and

relative IgA deficiency therefore appears to be related to the particular IgA variant(s) that are present within this breed.

In the present study, only variant C was found in all of the GSD tested (i.e. all GSD were homozygous for variant C). This suggests that the presence of variant C allele predisposes German Shepherd dogs to gastrointestinal and other IgA-related disorders known to be more prevalent in GSDs than in other breeds. Variant C may also result in a deficiency of IgA, possibly by making the molecule more susceptible to degradation. Furthermore, tests on a panel of non-specific dietary sensitive dogs showed that 80% had one or more variant C allele, in comparison to 39% of dogs in a random population. This suggests that the presence of a variant C allele increases susceptibility to gastrointestinal disease.



## SEQUENCE LISTING

<110> MARS, INCORPORATED  
 5 <120> ALLELIC VARIANTS  
 <130> N.89234 GCW  
 <160> 18  
 10 <170> PatentIn version 3.0  
 <210> 1  
 <211> 1611  
 15 <212> DNA  
 <213> Canis familiaris  
 <220>  
 <221> CDS  
 20 <222> (3)..(305)  
 <220>  
 <221> variation  
 <222> (514)..(519)  
 25 <223> n indicates variable number of CT repeats  
 <220>  
 <221> CDS  
 <222> (557)..(575)  
 30 <220>  
 <221> CDS  
 <222> (579)..(897)  
 35 <220>  
 <221> CDS  
 <222> (1099)..(1486)  
 40 <400> 1  
 ag tcc aaa acc agc ccc agt gtg ttc ccg ctg agc ctc tgc cac cag 47  
 Ser Lys Thr Ser Pro Ser Val Phe Pro Leu Ser Leu Cys His Gln  
 1 5 10 15  
 gag tca gaa ggg tac gtg gtc atc ggc tgc ctg gtg cag gga ttc ttc 95  
 Glu Ser Glu Gly Tyr Val Val Ile Gly Cys Leu Val Gln Gly Phe Phe  
 20 25 30  
 cca ccg gag cct gtg aac gtg acc tgg aat gcc ggc aag gac agc aca 143  
 Pro Pro Glu Pro Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr  
 35 40 45  
 tct gtc aag aac ttc ccc ccc atg aag gct gct acc gga agc cta tac 191  
 Ser Val Lys Asn Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr  
 50 55 60  
 acc atg agc agc cag ttg acc ctg cca gcc gcc cag tgc cct gat gac 239  
 Thr Met Ser Ser Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp  
 65 70 75  
 tcg tct gtg aaa tgc caa gtg cag cat gct tcc agc ccc agc aag gca 287  
 Ser Ser Val Lys Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala  
 80 85 90 95  
 gtg tct gtg ccc tgc aaa ggtcagaggg caggctgggg aggggcaggg 335  
 Val Ser Val Pro Cys Lys  
 100  
 gccccacatc ctactctga ccctccactt ggagttctgg cccaaggac actccacggg 395

gagcacagtg ggctgctggg ctgagctccc agcaagtggc caaggtggg cctccatgaa 455  
 ggacctggag ggtggcaggg ggcaggcagg cagaggggtgc aactgacct gttccaatnn 515  
 nnnnctctct ctctctctct ctctctctct ctgctcctga a gat aac agt cat ccg 571  
 Asp Asn Ser His Pro 105  
 tgt c nnn at cca tgt ccc tcg tgc aat gag ccc cgc ctg tca cta cag 619  
 Cys His Pro Cys Pro Ser Cys Asn Glu Pro Arg Leu Ser Leu Gln 110 115 120  
 aag cca gcc ctc gag gat ctg ctt tta ggc tcc aat gcc agc ctc aca 667  
 Lys Pro Ala Leu Glu Asp Leu Leu Glu Gly Ser Asn Ala Ser Leu Thr 125 130 135  
 tgc aca ctg agt ggc ctg aaa gac ccc aag ggt gcc acc ttc acc tgg 715  
 Cys Thr Leu Ser Gly Leu Lys Asp Pro Lys Gly Ala Thr Phe Thr Trp 140 145 150  
 aac ccc tcc aaa ggg aag gaa ccc atc cag aag aat cct gag cgt gac 763  
 Asn Pro Ser Lys Gly Lys Glu Pro Ile Gln Lys Asn Pro Glu Arg Asp 155 160 165  
 tcc tgt ggc tgc tac agt gtg tcc agt gtc cta cca ggc tgt gct gat 811  
 Ser Cys Gly Cys Tyr Ser Val Ser Ser Val Leu Pro Gly Cys Ala Asp 170 175 180 185  
 cca tgg aac cat ggg gac acc ttc tcc tgc aca gcc acc cac cct gaa 859  
 Pro Trp Asn His Gly Asp Thr Phe Ser Cys Thr Ala Thr His Pro Glu 190 195 200  
 tcc aag agc ccg atc act gtc agc atc acc aaa acc ac aggtgggccc 907  
 Ser Lys Ser Pro Ile Thr Val Ser Ile Thr Lys Thr Thr 205 210  
 agaccctgcc cgtgaggcac tgcttggcac acaaaagttt gtgaggcaac tcctaagcct 967  
 gcttccttcc tctagcccct gggcttgggt gctcccaccc acattttaca aagggaaact 1027  
 gtggcatggg gtgctatggg gaagaaggct cttccccac cccagatccc tgacctggct 1087  
 ctctgtcctg c a gag cac atc ccg ccc cag gtc cac ctg ctg ccg ccg 1135  
 Glu His Ile Pro Pro Gln Val His Leu Leu Pro Pro 215 220 225  
 ccg tcg gaa gag ctg gcc ctc aat gag ctg gtg aca ctg acg tgc ttg 1183  
 Pro Ser Glu Glu Leu Ala Leu Asn Glu Leu Val Thr Leu Thr Cys Leu 230 235 240  
 gtg agg ggc ttc aaa cca aaa gat gtg ctc gta cga tgg ctg caa ggg 1231  
 Val Arg Gly Phe Lys Pro Lys Asp Val Leu Val Arg Trp Leu Gln Gly 245 250 255  
 acc cag gag cta ccc caa gag aag tac ttg acc tgg gag ccc ctg aag 1279  
 Thr Gln Glu Leu Pro Gln Glu Lys Tyr Leu Thr Trp Glu Pro Leu Lys 260 265 270  
 gag cct gac cag acc aac atg ttt gcc gtg acc agc atg ctg agg gtg 1327  
 Glu Pro Asp Gln Thr Asn Met Phe Ala Val Thr Ser Met Leu Arg Val 275 280 285 290  
 aca gcc gaa gac tgg aag cag ggg gag aag ttc tcc tgc atg gtg ggc 1375  
 Thr Ala Glu Asp Trp Lys Gln Gly Glu Lys Phe Ser Cys Met Val Gly 295 300 305  
 cac gag gct ctg ccc atg tcc ttc acc cag aag acc atc gac cgc ctg 1423  
 His Glu Ala Leu Pro Met Ser Phe Thr Gln Lys Thr Ile Asp Arg Leu 310 315 320  
 gcg ggt aaa ccc acc cac gtc aac gtg tct gtg gtc atg gca gag gtg 1471

Ala Gly Lys Pro Thr His Val Asn Val Ser Val Val Met Ala Glu Val  
 325 330 335

5 gac ggc atc tgc tac taaaccgccc aatcttcctt ccctaaataa actccatgct 1526  
 Asp Gly Ile Cys Tyr 340

10 tgcccaaagc agccccgtgc ttccatcagg ccgcctgtct gtccatattc ggggtctgtg 1586  
 gcatactgag gcaggggtag agctc 1611

<210> 2  
 <211> 343  
 <212> PRT  
 <213> Canis familiaris

15 <400> 2  
 Ser Lys Thr Ser Pro Ser Val Phe Pro Leu Ser Leu Cys His Gln Glu  
 1 5 10 15

20 Ser Glu Gly Tyr Val Val Ile Gly Cys Leu Val Gln Gly Phe Phe Pro  
 20 25 30

25 Pro Glu Pro Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser  
 35 40 45

Val Lys Asn Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr  
 50 55 60

30 Met Ser Ser Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser  
 65 70 75 80

Ser Val Lys Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val  
 85 90 95

35 Ser Val Pro Cys Lys Asp Asn Ser His Pro Cys His Pro Cys Pro Ser  
 100 105 110

Cys Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp Leu  
 115 120 125

40 Leu Leu Gly Ser Asn Ala Ser Leu Thr Cys Thr Leu Ser Gly Leu Lys  
 130 135 140

45 Asp Pro Lys Gly Ala Thr Phe Thr Trp Asn Pro Ser Lys Gly Lys Glu  
 145 150 155 160

Pro Ile Gln Lys Asn Pro Glu Arg Asp Ser Cys Gly Cys Tyr Ser Val  
 165 170 175

50 Ser Ser Val Leu Pro Gly Cys Ala Asp Pro Trp Asn His Gly Asp Thr  
 180 185 190

Phe Ser Cys Thr Ala Thr His Pro Glu Ser Lys Ser Pro Ile Thr Val  
 195 200 205

55 Ser Ile Thr Lys Thr Thr Glu His Ile Pro Pro Gln Val His Leu Leu  
 210 215 220

60 Pro Pro Pro Ser Glu Glu Leu Ala Leu Asn Glu Leu Val Thr Leu Thr  
 225 230 235 240

Cys Leu Val Arg Gly Phe Lys Pro Lys Asp Val Leu Val Arg Trp Leu  
 245 250 255

65 Gln Gly Thr Gln Glu Leu Pro Gln Glu Lys Tyr Leu Thr Trp Glu Pro  
 260 265 270

Leu Lys Glu Pro Asp Gln Thr Asn Met Phe Ala Val Thr Ser Met Leu

275                      280                      285

Arg Val Thr Ala Glu Asp Trp Lys Gln Gly Glu Lys Phe Ser Cys Met  
290                      295                      300

5 Val Gly His Glu Ala Leu Pro Met Ser Phe Thr Gln Lys Thr Ile Asp  
305                      310                      315                      320

10 Arg Leu Ala Gly Lys Pro Thr His Val Asn Val Ser Val Val Met Ala  
325                      330                      335

Glu Val Asp Gly Ile Cys Tyr  
340

15 <210> 3  
<211> 542  
<212> DNA  
<213> Canis familiaris

20 <220>  
<221> CDS  
<222> (2)..(199)

25 <220>  
<221> variation  
<222> (408)..(435)  
<223> variable number of CT repeats [CT]n  
n = 12 or 14

30 <220>  
<221> CDS  
<222> (445)..(540)

35 <400> 3  
t gtg aac gtg acc tgg aat gcc ggc aag gac agc aca tct gtc aag aac 49  
Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn  
1                      5                      10                      15

40 ttc ccc ccc atg aag gct gct acc gga agc cta tac acc atg agc agc 97  
Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser  
20                      25                      30

cag ttg acc ctg cca gcc gcc cag tgc cct gat gac tcg tct gtg aaa 145  
Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys  
35                      40                      45

45 tgc caa gtg cag cat gct tcc agc ccc agc aag gca gtg tct gtg ccc 193  
Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro  
50                      55                      60

50 tgc aaa ggtcagaggg caggctgggg aggggcaggg gccccacatc ctcactctga 249  
Cys Lys  
65

55 ccctccactt ggagccgtgt ccccaaggac actccacggg gaggacagtg ggctgctggg 309  
ctgagctccc agcaagtggc caaggtgggg cctccatgaa ggacctggag ggtggcaggg 369  
ggcaggcagg cagaggtgac acactgacct gttccaatct ctctctctct ctctctctct 429

60 ctctctgctc ctgaa gat aac agt cat ccg tgt cat cca tgt ccc tcg tgc 480  
Asp Asn Ser His Pro Cys His Pro Cys Pro Ser Cys  
70                      75

aat gag ccc cgc ctg tca cta cag aag cca gcc ctc gag gat ctg ctt 528  
Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp Leu Leu  
80                      85                      90

65 tta ggc tcc aat gc 542  
Leu Gly Ser Asn

95

<210> 4  
 <211> 98  
 <212> PRT  
 <213> Canis familiaris

<400> 4  
 Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn  
 1 5 10 15

Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser  
 20 25 30

Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys  
 35 40 45

Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro  
 50 55 60

Cys Lys Asp Asn Ser His Pro Cys His Pro Cys Pro Ser Cys Asn Glu  
 65 70 75 80

Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp Leu Leu Leu Gly  
 85 90 95

Ser Asn

<210> 5  
 <211> 520  
 <212> DNA  
 <213> Canis familiaris

<220>  
 <221> CDS  
 <222> (2)..(199)

<220>  
 <221> mutation  
 <222> (73)..(73)  
 <223> C or T

<220>  
 <221> mutation  
 <222> (118)..(118)  
 <223> C or T

<220>  
 <221> variation  
 <222> (408)..(419)  
 <223> variable number of CT repeats [CT]n  
 n = 6 or 11

<220>  
 <221> CDS  
 <222> (429)..(518)

<400> 5  
 t gtg aac gtg acc tgg aat gcc ggc aag gac agc aca tct gtc aag aac  
 Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn  
 1 5 10 15

ttc ccc ccc atg aag gct gct act gga agc cta tac acc atg agc agc  
 Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser  
 20 25 30

cag ttg acc ctg cca gct gcc cag tgc cct gat gac tcg tct gtg aaa

49

97

145

Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys  
 35 40 45

5 tgc caa gtg cag cat gct tcc agc ccc agc aag gca gtg tct gtg ccc 193  
 Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro  
 50 55 60

10 tgc aaa ggtcagaggg caggctgggg aggggcaggg gccccacatc ctactctga 249  
 Cys Lys  
 65

15 ccctccactt ggagccgtgt cccaaggac actccacggg gaggacagtg ggtgctggg 309  
 ctgagctccc agcaagtggc caaggtgggg cctccatgaa ggacctggag ggtggcaggg 369  
 ggcaggcagg cagaggtgac acactgacct gttccaatct ctctctctct gctcctgaa 428

gat aac tgt cct cat ctg tgt ccc tca tgc aat gag ccc cac ctg tca 476  
 Asp Asn Cys Pro His Leu Cys Pro Ser Cys Asn Glu Pro His Leu Ser  
 70 75 80

20 cta cag aag cca gcc ctc gag gat ctg ctt tta ggc tcc aat gc 520  
 Leu Gln Lys Pro Ala Leu Glu Asp Leu Leu Leu Gly Ser Asn  
 85 90 95

25 <210> 6  
 <211> 96  
 <212> PRT  
 <213> Canis familiaris

30 <400> 6  
 Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn  
 1 5 10 15

35 Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser  
 20 25 30

Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys  
 35 40 45

40 Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro  
 50 55 60

Cys Lys Asp Asn Cys Pro His Leu Cys Pro Ser Cys Asn Glu Pro His  
 65 70 75 80

45 Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp Leu Leu Leu Gly Ser Asn  
 85 90 95

50 <210> 7  
 <211> 548  
 <212> DNA  
 <213> Canis familiaris

55 <220>  
 <221> CDS  
 <222> (2)..(199)

60 <220>  
 <221> mutation  
 <222> (118)..(118)  
 <223> C or T

65 <220>  
 <221> CDS  
 <222> (442)..(546)

<400> 7  
 t gtg aac gtg acc tgg aat gcc ggc aag gac agc aca tct gtc aag aac 49  
 Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn

	1	5	10	15	
5	ttc ccc ccc atg aag gct gct act gga agc cta tac acc atg agc agc				97
	Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser	20	25	30	
10	cag ttg acc ctg cca gcc gcc cag tgc cct gat gac tcg tct gtg aaa				145
	Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys	35	40	45	
15	tgc caa gtg cag cat gct tcc agc ccc agc aag gca gtg tct gtg ccc				193
	Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro	50	55	60	
20	tgc aaa ggtcagaggg caggctgggg aggggcaggg gcccacatc ctctctctga				249
	Cys Lys	65			
25	ccctccactt ggagccgtgt cccaaggac actccacggg gaggacagtg ggctgctggg				309
	ctgagctccc agcaagtggc caaggtgggg cctccatgaa ggacctggag ggtggcaggg				369
	ggcaggcagg cagaggggtgc acactgacct gttccaatct ctctctctct ctctctctct				429
30	ctctctctct ca gct cct gaa gat aac tgt cat ccg tgt cat cca tgt ccc				480
	Ala Pro Glu Asp Asn Cys His Pro Cys His Pro Cys Pro	70	75		
35	tcg tgc aat gag ccc cgc ctg tca cta cag aag cca gcc ctc gag gat				528
	Ser Cys Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp	80	85	90	95
40	ctg ctt tta ggc tcc aat gc				548
	Leu Leu Leu Gly Ser Asn	100			
45	<210> 8				
	<211> 101				
	<212> PRT				
	<213> Canis familiaris				
50	<400> 8				
	Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn	1	5	10	15
55	Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser	20	25	30	
	Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys	35	40	45	
60	Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro	50	55	60	
	Cys Lys Ala Pro Glu Asp Asn Cys His Pro Cys His Pro Cys Pro Ser	65	70	75	80
65	Cys Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp Leu	85	90	95	
	Leu Leu Gly Ser Asn	100			
	<210> 9				
	<211> 545				
	<212> DNA				
	<213> Canis familiaris				
	<220>				
	<221> CDS				

<222> (2)..(199)

<220>

<221> CDS

5 <222> (436)..(543)

<400> 9

t gtg aac gtg acc tgg aat gcc ggc aag gac agc aca tct gtc aag aac 49

Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn

10 1 5 10 15

ttc ccc ccc atg aag gct gct acc gga agc cta tac acc atg agc agc 97

Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser

15 20 25 30

cag ttg acc ctg cca gcc gcc cag tgc cct gat gac tcg tct gtg aaa 145

Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys

35 40 45

tgc caa gtg cag cat gct tcc agc ccc agc aag gca gtg tct gtg ccc 193

Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro

50 55 60

tgc aaa ggtcagaggg caggctgggg aggggcaggg gccccacatc ctactctga 249

Cys Lys

65

ccctccactt ggagcctgt cccaaggac actccacggg gaggacagtg ggctgctggg 309

ctgagctccc agcaagtggc caaggtgggg cctccatgaa ggacctggag ggtggcaggg 369

30 ggcaggcagg cagagggtgc acactgacct gtccaatct ctctctctct ctctctctct 429

ctctca gct cct gaa gat aac tgt cat ccg tgt cct cat cca agt ccc 477

Ala Pro Glu Asp Asn Cys His Pro Cys Pro His Pro Ser Pro

70 75 80

35 tcg tgc aat gag ccc cgc ctg tca cta cag aag cca gcc ctc gag gat 525

Ser Cys Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp

85 90 95

40 ctg ctt tta ggc tcc aat gc 545

Leu Leu Leu Gly Ser Asn

100

<210> 10

45 <211> 102

<212> PRT

<213> Canis familiaris

<400> 10

50 Val Asn Val Thr Trp Asn Ala Gly Lys Asp Ser Thr Ser Val Lys Asn

1 5 10 15

Phe Pro Pro Met Lys Ala Ala Thr Gly Ser Leu Tyr Thr Met Ser Ser

20 25 30

55 Gln Leu Thr Leu Pro Ala Ala Gln Cys Pro Asp Asp Ser Ser Val Lys

35 40 45

Cys Gln Val Gln His Ala Ser Ser Pro Ser Lys Ala Val Ser Val Pro

50 55 60

60 Cys Lys Ala Pro Glu Asp Asn Cys His Pro Cys Pro His Pro Ser Pro

65 70 75 80

65 Ser Cys Asn Glu Pro Arg Leu Ser Leu Gln Lys Pro Ala Leu Glu Asp

85 90 95

Leu Leu Leu Gly Ser Asn



100

5     <210> 11  
       <211> 20  
       <212> DNA  
       <213> Artificial  
  
       <220>  
       <223> Primer  
 10     <400> 11  
       tgtgaacgtg acctggaatg 20  
  
 15     <210> 12  
       <211> 20  
       <212> DNA  
       <213> Artificial  
  
 20     <220>  
       <223> Primer  
  
       <400> 12 20  
       gcattggagc ctaaaagcag  
 25  
  
       <210> 13  
       <211> 20  
       <212> DNA  
 30     <213> Artificial  
  
       <220>  
       <223> Primer  
  
 35     <400> 13 20  
       tgtgccctgc aaagataaca  
  
       <210> 14  
 40     <211> 20  
       <212> DNA  
       <213> Artificial  
  
       <220>  
 45     <223> Primer  
  
       <400> 14 20  
       agggctggct tctgtagtga  
  
 50     <210> 15  
       <211> 20  
       <212> DNA  
       <213> Artificial  
  
 55     <220>  
       <223> Primer  
  
       <400> 15 20  
 60     cgtctgtgaa atgccaagtg  
  
       <210> 16  
 65     <211> 20  
       <212> DNA  
       <213> Artificial

<220>  
<223> Primer

5 <400> 16  
agggctggct tctgtagtga

20

10 <210> 17  
<211> 26  
<212> DNA  
<213> Artificial

15 <220>  
<223> Probe  
  
<400> 17  
gtcatccatg tccctcgtgc aatgag

26

20 <210> 18  
<211> 1789  
<212> DNA  
<213> Canis familiaris

25 <400> 18  
agtgaacctag cgtgtcattc tgacccagggt ctccggcatat gaactgcatg accttgggct 60  
gtcactgacc atctctatgc agtttctctt agtgcaaaaga aaaaatagcc ctcaccctgc 120  
ctgtgaggcc atgtaagggg tccagacagc actggccccc cagctcacag agtgcctgt 180  
gtcacagagt ccaaaaccag cccagtggtg ttcccgtga gcctctgcca ccaggagtca 240  
30 gaagggtacg tggtcacggt ctgcctgggt cagggtattct tcccaccgga gcctgtgaac 300  
gtgacctgga atgcccggcaa ggacagcaca tctgtcaaga acttcccccc catgaaggct 360  
gctaccggaa gcctatacac catgagcagc cagttgaccc tgccagccgc ccagtgcct 420  
gatgactcgt ctgtgaaatg ccaagtgcag catgcttcca gcccagcaa ggagtgctct 480  
35 gtgccctgca aaggctcagag ggcaggctgg ggaggggcag gggccccaca tctcactct 540  
gacctccac ttggagttct ggcccgaagg aactccacg gggaggacag tgggctgctg 600  
ggctgagctc ccagcaagtg gccaaagtg ggctccatg aaggacctg aggggtggcag 660  
ggggcaggca ggcagagggt gcacactgac ctgttccaat ctctctctct ctctctctct 720  
40 ctotctctgc tcctgaagat aacagtcac cgtgtcatcc atgtccctcg tgcaatgagc 780  
ccgcctgtc actacagaag ccagccctcg aggatctgct tttaggctcc aatgccagcc 840  
tcacatgcac actgagtggt ctgaaagacc ccaagggtgc caccttacc tggaacccct 900  
ccaaagggaa ggaaccatc cagaagaatc ctgagcgtga ctctgtggc tgctacagtg 960  
tgtccagtgt cctaccaggc tgtgctgac gatggaacca tggggacacc ttctcctgca 1020  
cagccacca cctgaatcc aagagccga tcaactgtcag catcacaaa accacagggtg 1080  
45 ggcccagacc ctgcccgtga ggcactgctt ggcacacaaa agtttgtgag gcaactccta 1140  
agcctgtctc ctctctctag cccctgggct tgggtgtcc cacccacatt ttacaaaggg 1200  
aaactgtggc atgggggtgt atggggaaga aggtctctcc cccaccccag atccctgacc 1260  
tggctctctg tcctgcagag cacatccgc cccaggtcca cctgctgccg ccgcccgtcg 1320  
aagagctggc cctcaatgag ctggtgacac tgacgtgctt ggtgaggggc ttcaaaccac 1380  
50 aagatgtgtc cgtacgatgg ctgcaaggga cccaggagct accccaagag aagtacttga 1440  
cctgggagcc cctgaaggag cctgaccaga ccaacatgtt tgccgtgacc agcatgctga 1500  
gggtgacagc cgaagactgg aagcaggggg agaagttctc ctgcatggtg ggccacgagg 1560  
ctctgccat gtccttcacc cagaagacca tcgaccgcct ggcgggtaaa cccaccacg 1620  
tcaacgtgtc tgtggtcatg gcagaggtgg acggcatctg ctactaaacc gcccaatctt 1680  
55 cctccctaa ataaactcca tgcttgcca aagcagcccc gtgcttccat caggccgcct 1740  
gtctgtccat attcgggggtc tgtggcatat tgaggcaggg gtagagctc 1789

CLAIMS

1. A method for determining susceptibility to an IgA-related disorder in an animal, the method comprising:

5 a) identifying the or each IgA allelic variant present in a sample from the animal; and  
b) thereby determining whether the animal is susceptible to an IgA-related disorder.

2. A method according to claim 1, wherein the animal is a dog.

10 3. A method according to claim 1 or 2, wherein the IgA-related disorder is gastrointestinal, skin, respiratory, rheumatoid or periodontal disease.

4. A method according to claim 3, wherein the disease is diarrhoea, small intestinal bacterial overgrowth, inflammatory bowel disease, perianal fistulas, atopic dermatitis, pyoderma, anal furunculosis, malassezia infestans or disseminated  
15 aspergillosis.

5. A method according to any one of the preceding claims, wherein identification of the allelic variant comprises detecting one or more polymorphisms in the hinge region of the IgA allelic variant, or a polymorphism which is in linkage disequilibrium with such a polymorphism.

20 6. A method according to claim 5, wherein the polymorphism is at any one of the following positions in relation to SEQ ID NO: 1:

- position 179 [C/T];
- position 370 [T/C];
- position 371 [T/C];
- 25 - position 372 [C/G];
- position 375 [G/T];
- positions 514 to 546 [number of CT repeats];
- position 547 [T/A];
- position 563 [A/T];
- 30 - positions 563 to 571 [deletion];
- positions 576 to 578 [addition];
- position 582 [C/T];
- position 583 [A/G];
- position 584 [T/A];

- position 592 [G/A]; or
- position 606 [G/A];

7. A method according to any one of the preceding claims, wherein the presence of at least one variant C allele indicates susceptibility to an IgA-related disorder.

8. A method according to any one of the preceding claims, wherein step (a) comprises contacting a polynucleotide encoding an IgA allelic variant with a specific binding agent for an allelic variant and determining whether the agent binds to the polynucleotide, wherein binding of the agent to the polynucleotide indicates the presence of the allelic variant.

9. A method according to claim 8 wherein the agent is a polynucleotide which is able to bind a polynucleotide encoding the IgA allelic variant but which does not bind a polynucleotide encoding a different IgA allelic variant.

10. A method according to any one of claims 1 to 7, wherein step (a) comprises contacting an IgA allelic variant polypeptide with a specific binding agent for an allelic variant and determining whether the agent binds to the polypeptide, wherein binding of the agent to the polypeptide indicates the presence of the allelic variant.

11. A method according to any one of claims 1 to 7 wherein the allelic variant is detected by measuring the mobility of an IgA allelic variant polypeptide or a polynucleotide encoding an IgA allelic variant during gel electrophoresis.

12. A probe, primer or antibody which is capable of detecting an IgA allelic variant.

13. A kit for carrying out the method of any one of claims 1 to 11 comprising means for detecting an IgA allelic variant.

14. A kit according to claim 13, comprising a probe, primer or antibody according to claim 12.

15. A method of preparing customised food for an animal which is susceptible to an IgA-related disorder, the method comprising:

- (a) determining whether the animal is susceptible to an IgA-related disorder by a method according to any one of claims 1 to 11; and
- (b) preparing food suitable for the animal.

16. A method according to claim 15, wherein the customised animal food comprises ingredients which prevent or alleviate an IgA-related disorder, and/or does not comprise ingredients which contribute to or aggravate an IgA-related disorder.

17. A method according to claim 16 wherein the customised animal food comprises  $\beta$ -glucans, glutamine, probiotics, oligosaccharides, exogenous IgA, hypoallergenic protein, hydrolysed protein, vitamin C, taurine, curcumin or aloe vera.

18. A method according to any one of claims 15 to 17, further comprising providing the food to the animal, the animal's owner or the person responsible for feeding the animal.

19. A method of providing a customised animal food, comprising providing food suitable for an animal which is susceptible to an IgA-related disorder to the animal, the animal's owner or the person responsible for feeding the animal, wherein the animal has been genetically determined to be susceptible to an IgA-related disorder.

20. A method for identifying an agent for the treatment of an IgA-related disorder, the method comprising:

(a) contacting an IgA allelic variant polypeptide or a polynucleotide which encodes an IgA allelic variant with a test agent; and

(b) determining whether the agent is capable of binding to the polypeptide or modulating the activity or expression of the polypeptide or polynucleotide.

21. Use of a compound which is therapeutic for an IgA-related disorder in the manufacture of a medicament for the prevention or treatment of an IgA-related disorder in an animal that has been identified as being susceptible to an IgA-related disorder by a method according to any one of claims 1 to 11.

22. A method of treating an animal for an IgA-related disorder, the method comprising administering to the animal an effective amount of a therapeutic compound which prevents or treats the disorder, wherein the animal has been identified as being susceptible to an IgA-related disorder by a method according to any one of claims 1 to 11.

23. A database comprising information relating to IgA allelic variants and optionally their association with IgA-related disorder(s).

24. A method for determining whether an animal is susceptible to an IgA-related disorder, the method comprising:

(a) inputting data of one or more IgA allelic variant(s) present in the animal to a computer system;

(b) comparing the data to a computer database, which database comprises information relating to IgA allelic variants and the IgA-related disorder susceptibility associated with the variants; and

(c) determining on the basis of the comparison whether the animal is susceptible to an IgA-related disorder.

25. A method according to claim 24, wherein the IgA allelic variant(s) are as defined in claim 6 or 7.

5 26. A computer program comprising program code means for performing all the steps of claim 24 when said program is run on a computer.

27. A computer program product comprising program code means stored on a computer readable medium for performing the method of claim 24 when said program product is run on a computer.

10 28. A computer program product comprising program code means on a carrier wave, which program code means, when executed on a computer system, instruct the computer system to perform a method according to claim 24.

29. A computer system arranged to perform a method according to claim 24 comprising:

15 (a) means for receiving data of the one or more IgA allelic variant(s) present in the animal;

(b) a module for comparing the data with a database comprising information relating to IgA allelic variants and the IgA-related disorder susceptibility associated with the variants; and

20 (c) means for determining on the basis of said comparison whether the animal is susceptible to an IgA-related disorder.

30. A method of preparing customised food for an animal which is susceptible to an IgA-related disorder, the method comprising:

25 (a) determining whether the animal is susceptible to an IgA-related disorder by a method according to claim 24;

(b) electronically generating a customised animal food formulation suitable for the animal;

30 (c) generating electronic manufacturing instructions to control the operation of food manufacturing apparatus in accordance with the customised animal food formulation; and

(d) manufacturing the customised animal food according to the electronic manufacturing instructions.

31. A computer system according to claim 29, further comprising:

(d) means for electronically generating a customised animal food formulation suitable for the animal;

(e) means for generating electronic manufacturing instructions to control the operation of food manufacturing apparatus in accordance with the customised animal food formulation; and

(f) a food product manufacturing apparatus.

32. Use of a computer system as defined in claim 31 to make a customised animal food product.

33. An isolated polynucleotide comprising:

10 (a) an IgA variant sequence that differs to SEQ ID NO: 1 at one or more polymorphic positions as defined in claim 6;

(b) any one of SEQ ID NO:s 3, 5, 7 or 9;

(c) a sequence that is complementary or is degenerate as a result of the genetic code to a sequence as defined in (a) or (b); or

15 (d) a fragment of (a), (b) or (c) which differs to SEQ ID NO: 1 at one or more polymorphic positions as defined in claim 6 and which is at least 10 nucleotides in length.

34. A polypeptide comprising:

(a) a sequence encoded by a polynucleotide according to claim 33;

20 (b) any one of SEQ ID NO:s 4, 6, 8 or 10; or

(c) a fragment of (a) or (b) which differs to SEQ ID NO: 2 at one or more polymorphic positions as defined in claim 6 and which is at least 10 amino acids in length.

ABSTRACTALLELIC VARIANTS

5       The invention relates to a method for determining susceptibility to an IgA-related disorder in an animal, the method comprising:


      a)     identifying the or each IgA allelic variant present in a sample from the animal; and

      b)     thereby determining whether the animal is susceptible to an IgA-related  
10   disorder.

15



Figure 1



B

A

Figure 2




Figure 3




Figure 4




Figure 5





Figure 6



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: \_\_\_\_\_**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**